FAO SPECIFICATIONS FOR PLANT PROTECTION PRODUCTS

CHLOROTOLURON

3-(3-chloro-p-tolyl)-1,1-dimethylurea

AND

FLUOMETURON

1,1-dimethyl-3-([a],[a],[a]-trifluoro-m-tolyl-urea

FOOD AND AGRICULTURE ORGANIZATION OF THE UNITED NATIONS Rome, 1990

Group on Pesticide Specifications

FAO Panel of Experts on Pesticide Specifications, Registration Requirements and Application Standards

Technical Secretary: Dr. A.V. Adam
Plant Protection Service
Plant Production and Protection Division

FAO

Via delle Terme di Caracalla 00100 Rome, Italy - Telex: 610181 FAO I

CONTENTS

DISCLAIMER	4
INTRODUCTION TO FAO SPECIFICATIONS DEVELOPED UNDER THE OLD PR	OCEDURE5
SUBMISSION OF DRAFT SPECIFICATIONS TO FAO	8
CHLOROTOLURON INFORMATION	9
CHLOROTOLURON TECHNICAL	10
CHLOROTOLURON WETTABLE POWDERS	11
CHLOROTOLURON WETTABLE POWDERS	12
CHLOROTOLURON AQUEOUS SUSPENSION CONCENTRATES	14
NOTES	16
FLUOMETURON INFORMATION	17
FLUOMETURON TECHNICAL	18
FLUOMETURON WETTABLE POWDERS	19
FLUOMETURON AQUEOUS SUSPENSION CONCENTRATES	21
NOTES	23

DISCLAIMER 1

FAO specifications are developed with the basic objective of promoting, as far as practicable, the manufacture, distribution and use of pesticides that meet basic quality requirements.

Compliance with the specifications does not constitute an endorsement or warranty of the fitness of a particular pesticide for a particular purpose, including its suitability for the control of any given pest, or its suitability for use in a particular area. Owing to the complexity of the problems involved, the suitability of pesticides for a particular purpose and the content of the labelling instructions must be decided at the national or provincial level.

Furthermore, pesticides which are manufactured to comply with these specifications are not exempted from any safety regulation or other legal or administrative provision applicable to their manufacture, sale, transportation, storage, handling, preparation and/or use.

FAO disclaims any and all liability for any injury, death, loss, damage or other prejudice of any kind that may arise as a result of, or in connection with, the manufacture, sale, transportation, storage, handling, preparation and/or use of pesticides which are found, or are claimed, to have been manufactured to comply with these specifications.

Additionally, FAO wishes to alert users to the fact that improper storage, handling, preparation and/or use of pesticides can result in either a lowering or complete loss of safety and/or efficacy.

FAO is not responsible, and does not accept any liability, for the testing of pesticides for compliance with the specifications, nor for any methods recommended and/or used for testing compliance. As a result, FAO does not in any way warrant or represent that any pesticide claimed to comply with a FAO specification actually does so.

-

¹ This disclaimer applies to all specifications published by FAO.

INTRODUCTION TO FAO SPECIFICATIONS DEVELOPED UNDER THE OLD PROCEDURE

Between 1975 and 2000, FAO published booklets of specifications for technical materials and related formulations of plant protection products. Revisions of, and additions to, already published specifications will be issued when necessary. However, all changes and revisions of FAO specifications are now subject to the new procedure described in the Manual on the development and use of FAO and WHO Specifications for Plant Protection Products, FAO Plant Production and Protection Paper No. 173, Rome 2002 (Revised First available FAO home the Edition only on the page of Internet http://www.fao.org/ag/agp/agpp/pesticid/)

FAO specifications developed under the old procedure are based on the requirements defined in the Fourth Edition of the *Manual on the development and use of FAO specifications for plant protection products*, Plant Production and Protection Paper No. 128, Rome 1995.

This manual contained detailed definitions and other essential background information on basic procedures and technical principles adopted by the group on Pesticide Specifications of the FAO Panel of Experts on Pesticide Specifications, Registration Requirements, Application Standards and Prior Informed Consent, such as:

1. Classes of Specifications (page 10 of the Manual).

<u>FAO</u> (full) specifications (Code "S"). Specifications that have all necessary requirements together with CIPAC (full) methods, or other collaboratively studied (proven) methods. 2/ and 3/.

<u>FAO Provisional specifications [Code (S)]</u> are those for which more evidence of the necessary parameters is available and where some collaborative study of the methods of analysis has been carried out.

<u>FAO Tentative specifications (Code "ts")</u> are those which have been recommended by FAO as preliminary specifications which are based on minimum requirements. The methods of analysis cited are normally supplied by the manufacturer or may already have been published or be the subject of collaborative work.

Wherever possible, standards for apparatus and common names for pesticides are those approved by the International Standards Organization (IS0).

- 2. Expression of Active Ingredient Content (page 18 of the Manual)
- for solids, liquid technical materials, volatile liquids (of maximum boiling point 50° C) and viscous liquids (with minimum kinematic viscosity of 1 x 10^{-3} m² / s at 20° C) the FAO Specification shall be based on g/kg expression of content;

- for all other liquids the active ingredient content of the product shall be declared in terms of g/kg or g/l at 20°C. If the buyer requires both g/kg and g/l at 20°C, then, in case of dispute, the analytical results shall be calculated as g/kg.

3. Tolerance on Content (page 19 of the Manual).

A declared content o£ active ingredient must be included in all specifications, and one of the problems immediately arising is the level of tolerance acceptable above the nominal figures. The tolerance is influenced by (a) the reproducibility o£ the method of analysis, (b) the sampling error and (c) the manufacturing variance.

Allowable variations in analytical results (i.e., tolerances in content of active ingredient) with respect to specific pesticide consignments are intended to cover reasonable variations in content of active ingredient. For examples of such permitted tolerances, see the table on page 20 of the Manual.

4. Containers/Packaging (page 32 of the Manual).

Containers shall comply with pertinent national and international transport and safety regulations.

- <u>Technical material, dustable powders and granules</u>

Containers shall be suitable, clean, dry and as specified, and shall not adversely affect, or be affected by, the product/material, but shall adequately protect it against external conditions.

Wettable Powders

The product shall be packed in suitable, clean, dry containers as specified in the order. The container shall provide all necessary protection against compaction, atmospheric moisture, oxidation, loss by vaporization and/or contamination to ensure that the product suffers no deterioration under normal transit and storage conditions.

The product shall be protected by an adequate moisture barrier. This may be a suitable bag of polyethylene or alternative means of giving equal or better protection.

- Solutions and emulsifiable concentrates

Containers shall be lined, where necessary, with a suitable material, or the interior surfaces treated to prevent corrosion and/or deterioration of the contents.

Additional information should be given in all specifications where particular pesticides present problems in packaging.

5. Biological information (page 33 of the Manual).

- Phytotoxicity

No test can be specified to cover possible phytotoxicity of formulation to all crops. When a crop is not mentioned in the instructions for use, purchasers should check with the supplier that the material is suitable, always provided that such a use is not restricted or legally forbidden.

- Wetting of crops

The dilute spray should satisfactorily wet the leaves of the specified crops when used in accordance with the instructions. Test method MT 53.2, CIPAC 1, p. 965 may be useful.

1/ Should national pesticide specifications developed from these approved FAO specifications deviate fend them, the national Authority responsible for making such changes is requested to inform the FAO Plant Protection Service of the nature of and the reasons for the modifications.

2/ Methods or analysis and miscellaneous techniques referred to in these specifications have been developed and adopted by CIPAC (Collaborative International Pesticides Analytical Council Ltd.). See CIPAC Handbooks, 1 (1970), 1A (1980), 1B t1983), 1C (1985) and ID (1988), CIPAC Proceedings 1980 and 1981, obtainable from Black Bear Press Limited, King's Hedges Road, Cambridge CB4 2PQ, England. The page numbers of specific methods are given in brackets in the specifications. A copy of a method not yet published can be obtained from the FAO Plant Protection Service.

3/ Information on standard waters for laboratory evaluation of pesticidal formulations will be found in "CIPAC Monograph 1, Standard Waters and an FAO survey on Naturally Occurring Waters" (1972). Black Bear Press Limited, King's Hedges Road, Cambridge CB4 2PO, England.

SUBMISSION OF DRAFT SPECIFICATIONS TO FAO

Any organization, commercial firm or interested individual is encouraged to submit relevant specifications, or proposals for revision of existing specifications, for pesticide products for consideration and possible adoption by FAO. Correspondence should be addressed to the Pesticide Management Group, Plant Production and Protection Division, FAO, Via delle Terme di Caracalla, 00153, Rome, Italy.

General guidelines in preparing draft specifications are given in the *Manual on the development and use of FAO and WHO Specifications for Plant Protection Products*, FAO Plant Production and Protection Paper No. 173, Rome 2002 (Revised First Edition available only on the FAO home page of the Internet at: http://www.fao.org/ag/agp/agpp/pesticid/)

.

Specifications which are considered suitable for further processing are assigned priorities and circulated to appropriate organizations and specialists to comment. Comments, together with other relevant information, are then reviewed in detail by the Group on Specifications of the FAO Panel of Experts on Pesticide Specifications, Registration Requirements, Application Standards and Prior Informed Consent. The drafts are converted into FAO Provisional Specifications, or full FAO Specifications.

CHLOROTOLURON INFORMATION

COMMON NAME: Chlorotoluron (ISO)

EMPIRICAL FORMULA: C10H13ClN2O

RMM: 212.7

CAS REGISTRY NUMBER: 15545-48-9

CIPAC CODE NUMBER: 217

CHEMICAL NAME:

3-(3-chloro-p-tolyl)-1,1-dimethylurea (IUPAC) N'-(3-chloro-4-methylphenyl)-N,N-dimethylurea (CA)

CHLOROTOLURON TECHNICAL

FAO Specification 217/TC/S (1990)

.1 DESCRIPTION

The material shall consist of chlorotoluron together with related manufacturing impurities and shall be a white to slightly yellow powder free from visible extraneous matter and added modifying agents.

.2 ACTIVE INGREDIENT

.2.1 Identity test (CIPAC 1A, 217/1/M/1.2, p.1151)

Where the identity of the active ingredient is in doubt, then it shall comply with at least one additional test.

.2.2 Chlorotoluron (CIPAC 1A, 217/1/M/1.3, p.1151)

The chlorotoluron content shall be declared (not less than 975 g/kg) and, when determined, the content obtained shall not differ from that declared by more than +/- 20g.

.3 IMPURITIES (CIPAC 1A, 217/1/M/1.4, p.1154)

3-(3-chloro-4-tolyl)-1-methylurea: Maximum 8 g/kg 3-(4-tolyl)-1,1-dimethylurea: Maximum 8 g/kg

CHLOROTOLURON WETTABLE POWDERS

FAO Specification 217/WP/S (1990)

.1 DESCRIPTION

The material shall consist of a homogeneous mixture of technical chlorotoluron [complying with the requirements of FAO Specification 217/TC/s (1990)] together with filler(s) and any other necessary formulants. It shall be in the form of a fine powder free visible extraneous matter and hard lumps.

.2 ACTIVE INGREDIENT

.2.1 Identity test (CIPAC 1A, 217/3/M/1.2, p.1155)

Where the identity of the active ingredient is in doubt, then the isolated active ingredient shall comply with at least one additional test.

.2.2 Chlorotoluron (CIPAC 1A, 217/3/M/1.3, p.1155)

The chlorotoluron content shall be declared (g/kg) a determined, the content obtained shall not differ from that by more than the following amounts:

Declared content (g/kg)
Up to 500
Above 500
Permitted tolerance
+/- 5% of the declared content
+/- 25 g/kg

.3 PHYSICAL PROPERTIES

.3.1 pH range (MT 75.2, CIPAC 1A, p.1590)

Range: If required, 6 to 10

.3.2 Wet sieve test (MT 59.3, CIPAC 1, P.981)

Maximum: 2% retained on a 75 µm test sieve.

CHLOROTOLURON WETTABLE POWDERS

FAO Specification 217/WP/S (1990)

.1 DESCRIPTION

The material shall consist of a homogeneous mixture of technical chlorotoluron [complying with the requirements of FAO Specification 217/TC/S (1990)] together with filler(s) and any other necessary formulants. It shall be in the form of a fine powder free from vlsible extraneous matter and hard lumps.

.2 ACTIVE INGREDIENT

.2.1 Identity test (CIPAC 1A, 217/3/M/1.2, p.1155)

Where the identity of the active ingredient is in doubt, then the isolated active ingredient shall comply with at least one additional

.2.2 Chlorotoluron (CIPAC 1A, 217/3/MV1.3, p.1155)

The chlorotoluron content shall be declared (g/kg) and, when determined, the content obtained shall not differ from that declared by more than the following amounts:

Declared content
Up to 500

Above 500

Permitted tolerance
+/- 5% of the declared content
+/- 25% g/kg

.3 PHYSICAL PROPERTIES

.3.1 pH range (MT 75.2, CIPAC 1A, p.1590)

Range: If required, 6 to 10

.3.2 Wet sieve test (MT 59.3, CIPAC 1, P.981)

Maximum: 2% retained on a 75 µm test sieve.

.3.3 Suspensibility (CIPAC 1A, 217/3/M/1.7, p.1156) (Notes 1 and 2)

A minimum of 60% of the chlorotoluron content found under .2.2 shall be in suspension after 30 minutes in CIPAC Standard Water C.

Alternatively, if the buyer requires other CIPAC Standard Waters to be used, then this shall be specified when ordering.

.3.4 Persistent foam (MT 47, CIPAC 1, p.954)

Maximum: 25 ml after 1 minute.

.3.5 Wetting of the product (MT 53.3.1, CIPAC 1, p.967)

The product shall be completely wetted in 1 minute without swirling.

.4 STORAGE STABILITY

.4.1 Stability at 54 C (MT 46.1.1, CIPAC 1, p.951)

After storage at 54 + -2°C for 14 days, the product shall continue to comply with .2.2, .3.1, .3.2 and . 3.3.

Note 1 The product should be tested at the highest and lowest rates of use recommended by the supplier, provided this does not exceed the conditions given in the method.

Note 2 This test will normally be carried out after the heat stability test .4.1.

CHLOROTOLURON AQUEOUS SUSPENSION CONCENTRATES

FAO Specification 217/SC/S (1990)

.1 DESCRIPTION

The material shall consist of suspension of fine particles oftechnical chlorotoluron [complying with the requirement specifications 217/TC/S (1990)] in an aqueous phase, together with suitable formulants.

After gentle agitation, the material shall be homogeneous and suitable for dilution in water.

.2 ACTIVE INGREDIENT

.2.1 Identity test (217/3/M/1.2, CIPAC 1A, p.1155)

Where the identity of the active ingredient is in doubt, isolated active ingredient shall comply with at least one additional test.

.2.2 Chlorotoluron (217/3/M/1.3, CIPAC 1A, p.1155)

The chlorotoluron content shall be declared (g/kg or g/l at 20 C, Note 2) and, when determined, the content obtained shall nol from that declared by more than the following amounts:

Declared content (g/kg or g/l)	<u>Permitted tolerance</u>
250 up to 500	+/- 5% of the declared content
Above 500	+/- 25 g/kg or g/l

.3 PHYSICAL PROPERTIES

.3.1 Mass per millilitre at 20°C (MT 3.3, CIPAC 1C, p.2247)

If required, the mass per millilitre (g/ml) at 20°C shall be declared.

.3.2 pH range (MT 75.2, CIPAC 1A, p.1590)

Range: 6 to 8.5

.3.3 Pourability (MT 148, CIPAC 1C, p.2282)

Maximum = "rinsed residue:" 0.6%

.3.4 Spontaneity of dispersion (MT 160, CIPAC 1C, p.2291)

A minimum of 95% of the chlorotoluron content found under .2.2 shall be in suspension after 5 minutes in CIPAC Standard Water C.

Alternatively, if the buyer requires other CIPAC Standard Waters to be used, then this shall be specified when ordering.

.3.5 Suspensibility (MT 161, CIPAC 1C, p.2294) (Note 3)

A minimum of 70% of the chlorotoluron content found under .2.2 shall be in suspension after 30 minutes in CIPAC Standard Water C.

Alternatively, if the buyer requires other CIPAC Standard Waters to be used, then this shall be specified when ordering.

.3.6 Wet sieve test (MT 59.3, CIPAC 1, p.981)

Maximum: 1% of the product shall be retained on a 75 µm test sieve.

.3.7 Persistent foam (MT 47.2, CIPAC 1C, p.2249)

Maximum: 25 ml after 1 minute

.4 STORAGE STABILITY

.4.1 Stability at 0 C (MT 39.1, CIPAC 1, p.930)

After storage at 0 + 1°C for 7 days, the product shall continue to comply with .3.4, .3.5 and .3.6.

.4.2 Stability at 54 C (MT 46.1, CIPAC 1, p.951)

After storage at $54 + 2^{\circ}$ C for 14 days (Note 4), the product shall continue to comply with .2.2, .3.3 and .3.5, and if required with .3.2, .3.4 and .3.6.

Notes

Note 1 Before sampling to verify product quality, inspect commercial container carefully. On standing suspension concentrates usually develop a concentration gradient from the top to the bottom of the container. This may even result in the appearance of a clear liquid on the top and/or of sediment on the bottom. Therefore, before sampling, homogenize the product according to the instructions given by the manufacturer or, in the absence of such instructions, by gently shaking of the commercial container (for example, by inverting the closed container several times). After this procedure, the container should not contain a sticky layer of non-dispersed product at the bottom. A suitable and simple method of checking for a non-dispersed layer is by probing with a glass rod. All the physical chemical tests must be carried out on a laboratory taken after the recommended homogenization procedure.

Note 2 Unless homogenization is carried out carefully, it is possible for the sample to become aerated. This can lead to errors in the determination of the active ingredient content in g/l. It is preferable, therefore, to determine the content in g/kg and, if necessary, employ the density in g/ml ex Clause .3.1 to calculate the active ingredient content in g/l.

If the buyer requires both g/kg and g/l at 20°C, then in case of dispute, the analytical results shall be calculated as g/kg.

Note 3 This test will normally only be carried out after stability test .4.2.

Note 4 Unless other temperatures and/or times are specified.

FLUOMETURON INFORMATION

COMMON NAME: Fluometuron (ISO)

EMPIRICAL FORMULA: C10H11F3N2O

RMM: 232.2

CAS REGISTRY NUMBER: 2164-17-2

CIPAC CODE NUMBER: 159

CHEMICAL NAMES:

l,l-Dimethyl-3-([a],[a],[a] -trifluoro-m-tolyl)-urea (IUPAC) N,N-dimethyl-N'-[3-(trifluoromethyl)-phenyl]-urea (CA)

FLUOMETURON TECHNICAL FAO Specification 159/TC/S (1990)

1. **DESCRIPTION**

The material shall consist of fluometuron together with related manufacturing impurities and shall be a white to creamy crystalline powder free from visible extraneous matter and added modifying agents.

2. ACTIVE INGREDIENT

.2.1 Identity test (CIPAC 1B, 159/TC/M/2, p.1842)

Where the identity of the active ingredient is in doubt, then it shall comply with at least one additional test.

.2.2 Fluameturon (CIPAC 1B, 159/TC/M/3, p.1842)

The fluometuron content shall be declared (not less than 940 g/kg) and, when determined, the content obtained shall not differ from declared by more than ± -20 g.

FLUOMETURON WETTABLE POWDERS

FAO Specification 159/WP/S (1990)

.1 DESCRIPTION

The material shall consist of a homogeneous mixture of technical fluometuron [complying with the requirements of FAO Specification 159/TC/S (1990)] together with filler(s) and any other necessary formulants. It shall be in the form of a fine powder free from visible extraneous matter and hard lumps.

.2 ACTIVE INGREDIENT

.2.1 Identity test (CIPAC IB, 159/WP/M/2, p.1844)

Where the identity of the active ingredient is in doubt, then the isolated active ingredient shall comply with at least one additional test.

.2.2 Fluometuron (CIPAC 1B, 159/WP/M/3, p.1844)

The fluometuron content shall be declared (g/kg) and, when determined, the content obtained shall not differ from that declared by more than the following amounts:

<u>Declared content</u> <u>Permitted tolerance</u>

Up to 500 +/-5% of the declared content

Above 500 +/-25 g/kg

.3 PHYSICAL PROPERTIES

.3.1 pH range (MT 75.2, CIPAC 1A, p.1590)

Range: If required, 6 to 10

.3.2 Wet sieve test (MT 59.3, CIPAC 1, p.981)

Maximum: 2% retained on a 75 µm test sieve.

.3.3 Suspensibility (CIPAC) 1B, 159/WP/M/4, p.1844) (Notes 1 and 2)

A minimum of 60% of the fluometuron content found under .2.2 be in suspension after 30 minutes in CIPAC Standard Water C.

Alternatively, if the buyer requires other CIPAC Standard to be used, then this shall be specified when ordering.

.3.4 Persistent foam (MT 47, CIPAC 1, p.954)

Maximum: 25 ml after 1 minute.

.3.5 Wetting of the product (MT 53.3.1, CIPAC 1, p.967)

The product shall be completely wetted in 1 minute swirling.

.4 STORAGE STABILITY

.4.1 Stability at 54°C (MT 46.1.1, CIPAC 1, p.951)

After storage at 54 + -2°C for 14 days, the product shall to comply with .2.2, .3.1, .3.2 and .3.3.

Note 1 The product should be tested at the highest and lowest rates of use recommended by the supplier, provided this does not exceed the conditions given in the method.

Note 2 This test will normally be carried out after stability test .4.1.

FLUOMETURON AQUEOUS SUSPENSION CONCENTRATES

FAO Specification 159/SC/S (1990)

.1 DESCRIPTION

The material shall consist of a suspension of fine particles of technical fluometuron [complying with the requirement of FAO Specification 159/TC/S (1990)] in an aqueous phase, together with suitable formulants.

After gentle agitation, the material shall be homogeneous (Note 1) and suitable for dilution in water.

.2 ACTIVE INGREDIENT

.2.1 Identity test (159/TC/M/2, CIPAC 1B, p.1842)

Where the identity of the active ingredient is in doubt, then the isolated active ingredient shall comply with at least one additional test.

.2.2 Fluometuron (159/TC/M/3, CIPAC 1B, p.1842)

The fluometuron content shall be declared (g/kg or g/l at 20°C, Note 2) and, when determined, the content obtained shall not differ from that declared by more than the following amounts:

Declared content Permitted tolerance

250 up to 500 +/-5% of the declared content

Above 500 +/- 25 g/kg or g/l

.3 PHYSICAL PROPERTIES

.3.1 Mass per millilitre at 20 C (MT 3.3, CIPAC 1C, p.2247)

If required, the mass per millilitre (g/ml) at 20 C shall be declared.

.3.2 pH range (MT 75.2, CIPAC 1A, p.1590)

Range: 6 to 8.5

.3.3 Pourability (MT 148, CIPAC 1C, p.2282)

Maximum "rinsed residue:" 0.6%

.3.4 Spontaneity of dispersion (MT 160, CIPAC 1C, p.2291)

A minimum of 95% of the fluometuron content found under .2.2 shall be in suspension after 5 minutes in CIPAC Standard Water C.

Alternatively, if the buyer requires other CIPAC Standard Waters to be used, then this shall be specified when ordering.

.3.5 Suspensibility (MT 161, CIPAC 1C, p.2294) (Note 3)

A minimum of 70% of the fluometuron content found under .2.2 shall be in suspension after 30 minutes in CIPAC Standard Water C.

Alternatively, if the buyer requires other CIPAC Standard Waters to be used, then this shall be specified when ordering.

.3.6 Wet sieve test (MT 59.3, CIPAC 1, p.981)

Maximum: 1% of the product shall be retained on a 75 µm test sieve.

.3.7 Persistent foam (MT 47.2, CIPAC 1C, p.2249)

Maximum: 25 ml after 1 minute

.4 STORAGE STABILITY

.4.1 Stability at 0 C (MT 39.1, CIPAC 1, p.930)

After storage at 0 + -1°C for 7 days, the product shall continue to comply with .3.4, .3.5 and .3.6.

.4.2 Stability at 54 C (MT 46.1, CIPAC 1, p.951)

After storage at $54 + -2^{\circ}C$ for 14 days (Note 4), the produce shall continue to comply with .2.2, .3.3 and .3.5, and if require .3.2, .3.4 and .3.6.

Notes

Note 1 Before sampling to verify product quality, inspect commercial container carefully. On standing suspension concentrates usually develop a concentration gradient from the top to the bottom of the container. This may even result in the appearance of a clear liquid on the top and/or of sediment on the bottom. Therefore, before sampling, homogenize the product according to the instructions given by the manufacturer or, in the absence of such instructions, by gently shaking of the commercial container (for example, by inverting the closed container several times). After this procedure, the container should not contain a sticky layer of non-dispersed product at the bottom. A suitable and simple method of checking for a non-dispersed layer is by probing with a glass rod. All the physical chemical tests must be carried out on a laboratory taken after the recommended homogenization procedure.

Note 2 Unless homogenization is carried out carefully, it is possible for the sample to become aerated. This can lead to errors in the determination of the active ingredient content in g/l. It is preferable, therefore, to determine the content in g/kg and, if necessary, employ the density in g/ml ex Clause .3.1 to calculate the active ingredient content in g/l.

If the buyer requires both g/kg and g/l at 20°C, then in case of dispute, the analytical results shall be calculated as g/kg.

Note 3 This test will normally only be carried out after stability test .4.2.

Note 4 Unless other temperatures and/or times are specified.